

Inhibitory Neurophysiological Deficit as a Phenotype for Genetic Investigation of Schizophrenia

ROBERT FREEDMAN,* C.E. ADAMS, LAWRENCE E. ADLER, PAULA C. BICKFORD, JUDITH GAULT, JOSETTE G. HARRIS, HERBERT T. NAGAMOTO, ANN OLINCY, RANDAL G. ROSS, KAREN E. STEVENS, MERILYNE WALDO, AND SHERRY LEONARD

Many investigators have proposed that biological endophenotypes might facilitate the genetic analysis of schizophrenia. A deficit in the inhibition of the P50 evoked response to repeated auditory stimuli has been characterized as a neurobiological deficit in schizophrenia. This deficit is linked to a candidate gene locus, the locus of the $\alpha 7$ -nicotinic cholinergic receptor subunit gene on chromosome 15q14. Supportive evidence has been found by other investigators, including: 1) linkage of schizophrenia to the same locus; 2) linkage of bipolar disorder to the locus; and 3) replication of the existence of this neurobiological deficit and its relation to broader neuropsychological deficits in schizophrenia. It is certain that there are many genetic factors in schizophrenia and bipolar disorder; what is needed is a complete and precise description of the contribution of each individual factor to the pathophysiology of these illnesses. *J. Med. Genet. (Semin. Med. Genet.)* 97:58–64, 2000. © 2000 Wiley-Liss, Inc.

KEY WORDS: schizophrenia; genetics; nicotinic receptors; inhibition; auditory evoked potential

INTRODUCTION

A complex illness is one that does not result from a single genetic abnormality. Complexity is not only obvious from schizophrenia's non-Mendelian inheritance pattern, but it is also the likely situation for an illness that has no single defined biological deficit [Tsuang et al.,

1990]. After several major efforts, the current findings consist of many linkage signals of moderate effect on several different chromosomes [Cao et al., 1997; Blouin et al., 1998]. Similarly, despite intensive studies using a variety of different techniques, no single biological abnormality has emerged as the major

pathogenic factor. These two conundrums may well reflect the same problem, i.e., schizophrenia likely results from the interaction of several genetic and non-genetic pathogenic factors. A central thesis of our work has been that one solution to the dual genetic and biological complexity is to attempt to solve both of them simultaneously for discrete elements within schizophrenia itself. Linkage of a putative brain deficit in schizophrenia to a specific chromosomal site provides strong evidence that the deficit is indeed a discrete neurobiological dysfunction in the illness. Conversely, the most powerful phenotypes for linkage are likely to be discrete neurobiological deficits that result from the effect of a single gene. Although this strategy has not been widely used in schizophrenia, it has been successful in a number of other genetic illnesses, that, like schizophrenia, cluster in families but do not have Mendelian inheritance. Hemochromatosis is perhaps the best known; serum ferritin proved to be a better marker of genetic affection than the illness itself [Borecki et al., 1990]. Another example is colon cancer, where the expression of polyps, rather than clinical cancer itself, is the inherited phenotype [Leppert et al., 1990].

Robert Freedman is Professor of Psychiatry and Pharmacology at the University of Colorado and co-director of the Denver VA Medical Center Schizophrenia Research Center, where the other authors are also affiliated. He is a NARSAD distinguished investigator and a psychiatrist interested in the neurobiological basis of schizophrenia.

Cathy Adams, Assistant Professor of Psychiatry, is a neurobiologist interested in the neuroanatomy and neuroembryology of schizophrenia.

Lawrence E. Adler, Professor of Psychiatry, is a psychiatrist interested in the neuropharmacology of schizophrenia.

Paula C. Bickford, Associate Professor of Pharmacology, studies the neurobiology of sensory processing in animal models of schizophrenia and other illnesses.

Judith Gault, Instructor of Psychiatry, is a NARSAD young investigator and geneticist, who is interested in the genetic transmission of schizophrenia.

Josette Harris, Assistant Professor of Psychiatry, is a neuropsychologist interested in the inheritance of neuropsychological dysfunction in schizophrenia.

Ann Olincy, Assistant Professor of Psychiatry, is a NARSAD young investigator and a VA Research Career psychiatrist, who is interested in nicotinic receptor function in schizophrenia.

Randal G. Ross, Assistant Professor of Psychiatry, is a NARSAD mid career investigator and a VA Research Career psychiatrist, who is interested in children at risk for schizophrenia.

Karen Stevens, Assistant Professor of Psychiatry, is a neuropharmacologist who is interested in animal models of schizophrenia.

Merilyne Waldo, Assistant Professor of Psychiatry and NARSAD young investigator, is a psychologist interested in cross cultural studies of schizophrenia.

Sherry Leonard is Associate Professor of Psychiatry and VA Research Career Scientist and co-director of the VA Schizophrenia Center. She is interested in the molecular biology of nicotinic receptors and their relevance to the genetics of schizophrenia.

*Correspondence to: Professor of Psychiatry and Pharmacology, Vice-chairman, Department of Psychiatry, Campus Box C-268-71, University of Colorado Health Sciences Center, 4200 East Ninth Avenue, Denver CO 80262. E-mail: Robert.Freedman@uchsc.edu

THE SEARCH FOR A PHENOTYPE

Many investigators conceptualize schizophrenia as arising from more basic cognitive deficits. The biological bases of cognitive deficits in schizophrenia, however, are unknown and

One solution to the dual genetic and biological complexity is to attempt to solve both of them simultaneously.

are likely to be highly complex. Nearly every brain region, including the frontal and temporal lobes, the cerebellum, striatum, and thalamus, has been proposed to be involved. [Tamminga et al., 1992; Nelson et al., 1998; Andreasen et al., 1999;]. This panoply of biological abnormalities have made it difficult to assign specific deficits to specific causes. A possible simplification is to separate genetic from non-genetic influences. Therefore, several investigators, including ourselves, have attempted family studies, in which neuropsychological deficits have been isolated in members of the pedigree with varying degrees of genetic risk. Deficits in attentional function, for example, have been observed by us in parents of schizophrenics who have ancestral family histories of schizophrenia [Harris et al., 1996], and by others in children of schizophrenic mothers [Freedman et al., 1998], in siblings of schizophrenics [Cannon et al., 1994], and in clinically unaffected monozygotic twins of schizophrenics [Goldberg et al., 1990].

An even more elementary level of analysis is to consider biological deficits that might reflect deficits in a single neuronal mechanism. At this simpler level, there is the possibility of isolating a discrete deficit that might reflect the activity of a single gene. A number of biological deficits are shared by schizophrenics and their relatives. A variant of

segregation analysis was used to show that eye movement deficits are the phenotype for a single gene effect that also conveys risk for schizophrenia [Holzman et al., 1988], and linkage of eye movement deficits to the site of a previously known linkage for schizophrenia was demonstrated [Arolt et al., 1996]. Other deficits that are found in families include increased reaction time abnormalities [de Amicis et al., 1986], neuropsychological abnormalities [Cornblatt and Keilp, 1994], abnormalities in the P300 auditory evoked potential [Sham et al., 1994], and ventricular enlargement [Shihabuddin et al., 1996]. Our work on the P50 inhibitory deficit as an alternative phenotype is described below. To date, it is the only physiological abnormality whose linkage analysis has pointed to a specific candidate mechanism for neuronal dysfunction in schizophrenia [Freedman et al., 1997].

IDENTIFICATION OF AN INHIBITORY SENSORY GATING DYSFUNCTION IN SCHIZOPHRENICS AND THEIR RELATIVES

We applied the classical neurophysiological conditioning-testing paradigm, designed to study inhibition, to auditory evoked responses in schizophrenia [Adler et al., 1982]. Our goal was to identify an elementary neuronal function that could be studied in both humans and animal models. In this paradigm, pairs of identical stimuli, termed conditioning and test, are presented to the subject at various intrapair intervals. We discovered that at a range of intervals, including 500 msec, normals diminish the amplitude of the P50 wave response to the second or test stimulus, whereas schizophrenics have similar responses to both. Consistent differences in the inhibition of the P50 response between schizophrenics and normal subjects have been reported by us [Waldo et al., 1994; Griffith et al., 1995], by others to whom we introduced the paradigm [Ward et al., 1996], and by independent groups who have used similar or identical paradigms

[Boutros et al., 1991; Erwin et al., 1991; Judd et al., 1992; Jin et al., 1997; Clementz et al., 1998; Yee et al., 1998]. The inhibition is generally measured as the ratio of the amplitude of the test to the conditioning response. Figure 1 shows sample waveforms from four schizophrenic subjects and their parents. The P50 wave is identified and its amplitude measured relative to the preceding negativity by a computer algorithm [Nagamoto et al., 1989]. Families were chosen for this study to have apparent unilineal family histories of schizophrenia [Waldo et al., 1995]. In each case, the schizophrenic proband has a ratio of the amplitude of the test to the conditioning response greater than

A number of biological deficits are shared by schizophrenics and their relatives.

50%, that is generally the upper limit of normal values. This deficit in inhibition is shared by the parents who have an ancestral family history of schizophrenia, but not by the parents without such histories. A deficit in P50 inhibition correlates with neuropsychological evidence for a deficit in attention [Cullum et al., 1993; Erwin et al., 1998; Yee et al., 1998]. This feature of neuropsychological disability is also one of the familially transmitted aspects of schizophrenia [Goldberg et al., 1990; Cannon et al., 1994; Harris et al., 1996; Freedman et al., 1998].

BIOLOGICAL STUDIES POINT TO THE $\alpha 7$ -NICOTINIC RECEPTOR

The biological basis of the inhibitory abnormality was investigated in a parallel series of basic and clinical experiments. Briefly, the basic investigations showed that hippocampal pyramidal neurons are the source of the P20-N40 wave, the rodent equivalent of human P50 [Bickford-Wimer et al., 1990;




Figure 1. Evoked potential recordings from four nuclear families (A–D) with a schizophrenic offspring. The recordings show the conditioning and test responses. The P50 wave is indicated in each response by the computer generated tic below each wave and was measured relative to the preceding negativity indicated by the tic above. The percentage ratio of test to conditioning P50 amplitude is shown for each subject [from Waldo et al., 1995].

Miller et al., 1995]. The inhibition of response to the second stimulus is pre-synaptic, mediated by GABA_B receptors on glutamatergic afferents [Hershmann et al., 1995]. To release sufficient GABA release to activate these recep-

This deficit in inhibition is shared by the parents who have an ancestral family history of schizophrenia.

tors requires cholinergic activation of hippocampal interneurons via the reticula-septal pathway, mediated by $\alpha 7$ -

nicotinic receptors that we have shown to be localized on these interneurons [Freedman et al., 1993; Bickford et al., 1993; Miller et al., 1993, 1995; Breese et al., 1997; Frazier et al., 1998]. These receptors admit calcium to the interneurons, that produces nitric oxide, that in turn acts as a second messenger to prolong the effect of receptor activation on neuronal function [Adams and Stevens, 1998]. Loss of inhibition in animal models is produced by lesion of the cholinergic afferents, pharmacological blockade of the $\alpha 7$ -receptor, genetic deficiency in the $\alpha 7$ receptor in inbred mouse strains, or blockade of nitric oxide production by the $\alpha 7$ receptor [Luntz-Leybman et al., 1992; Bick-

ford et al., 1995; Stevens et al., 1996; Adams and Stevens et al., 1998].

Several clinical correlates of these basic science findings have been observed. First, $\alpha 7$ nicotinic receptor-mediated regulation of inhibition of auditory P50 response has been observed in schizophrenia. In schizophrenics and their relatives, nicotine transiently normalizes the deficit in P50 inhibition [Adler et al., 1992, 1993], as predicted from the animal model, with a pharmacology consistent with activation of the $\alpha 7$ receptor, i.e., failure of blockade by the muscarinic antagonists or the high affinity nicotinic receptor channel blocker mecamylamine [Freedman et al., 1994]. Furthermore, schizophrenics

can normalize not only this abnormality, but eye tracking abnormalities as well, at doses of nicotine that they achieve through their particularly heavy smoking [Olincy et al., 1997, 1998].

GENETIC LINKAGE OF PATHOPHYSIOLOGICAL DEFICITS IN SCHIZOPHRENIA ON CHROMOSOME 15

Several analyses of schizophrenics and their relatives characterized the heritability of the P50 inhibitory deficit [Waldo et al., 1994, 1995]. The finding of P50 inhibitory deficits in relatives of schizophrenics has been replicated by others [Clementz et al., 1997]. A preliminary genome-wide scan in multiplex schizophrenia families identified several areas that showed evidence for linkage of both the inhibitory deficit and schizophrenia, including chromosome 15q14 [Coon et al., 1993]. This area was subsequently discovered by another group to contain the $\alpha 7$ nicotinic receptor subunit gene [Chini et al., 1994]. Therefore, we isolated a new polymorphic marker (D15S1360) from a 180 kb YAC containing the $\alpha 7$ human gene. A second genome wide scan was performed in 9 multiplex families that yielded only one significant lod score ($P < 0.001$, verified by simulation), that exceeded 5.0 at D15S1360 without recombination for both two point and multipoint analyses. The lod score for schizophrenia was positive, but did not exceed 3.0. The difference reflects the reduced power of the schizophrenia phenotype, that was less than 50% penetrant. Thus, although a single allele of D15S1360 was generally co-inherited with both schizophrenia and the P50 inhibitory deficit, the number of persons with schizophrenia was less than the number with the P50 deficit, only some of whom were schizophrenic. Hence, the lod scores for the two traits were both positive, but of different magnitude [Freedman et al., 1997].

We have conducted additional studies in the NIMH Genetics Initiative

families to determine if there is further support for the initial linkage. Although these families have not been phenotyped for the P50 abnormality, they have been rigorously and independently diagnosed for schizophrenia. We found significant D15S1360 allele sharing among affected sib pairs ($P < 0.0024$) using SIBPAL [Leonard et al., 1998]. The NIMH group itself [Kaufmann et al., 1998] independently reported significant evidence for linkage in non-parametric analyses at ATC3C11, their only marker in the region. Two groups reported failures of replication at the marker we developed for this locus: one found an NPL score of 0.3 in schizophrenia and subsequently found linkage disequilibrium with the same marker in bipolar families [Neves-Pereira et al., 1998], and the other reported a lod score of 0.75 at the flanking highly polymorphic marker ACTC [Curtis et al., 1999]. Other groups have now shown inheritance at the 15q14 locus for bipolar disorder [Edenberg et al., 1997], one of them using markers specifically designed by us to examine the $\alpha 7$ nicotinic receptor locus [Neves-Pereira et al., 1999]. Several genetic loci have shown overlap between findings for bipolar disorder and schizophrenia, and the possibility that the two illnesses are part of the same spectrum has been raised by many investigators [Berrettini et al., 1997]. We have previously shown overlap between the P50 inhibitory deficits in bipolar disorder and schizophrenia in three studies [Adler et al., 1989, Baker et al., 1990]. Given the new genetic findings, further examination of the similarities and differences in phenotype associated with the $\alpha 7$ nicotinic receptor locus is warranted.

The $\alpha 7$ nicotinic receptor is thought to be a homopentamer in vitro; its in vivo structure is unknown. The gene is homologous across species and contains 10 exons. Humans, however, have a partially duplicated portion, that is about 1 million bases centromeric and thus within the linkage support region. It contains 4 novel exons, followed by exons 5–10 of the $\alpha 7$ gene [Gault et al., 1998]. Examination of the $\alpha 7$ gene and

its duplication to determine if there are indeed molecular alterations in the gene in schizophrenics is in progress.

BILINEAL INHERITANCE OF PHYSIOLOGICAL ABNORMALITIES IN CHILDHOOD ONSET SCHIZOPHRENIA

The inheritance of D15S1360 and the segregation of the P50 abnormality follow an apparent autosomal dominant pattern, so that most schizophrenics are expected to be heterozygous for any pathogenic allele. The data, however, do not exclude the possibility of co-dominance, a less common state in which individuals who have two pathogenic alleles have increased disability and, often, an earlier onset of illness. We hypothesized that childhood-onset schizophrenia might be the manifestation of co-dominance at the chromosome 15 locus. The severity of this illness initially resulted in it being considered a pan-developmental abnormality, perhaps unrelated to adult onset illness. More recently, phenotypic similarities between adult and childhood-onset illnesses have been emphasized [McKenna et al., 1994]. In many genetic illnesses, patients with increased severity and earlier onset have increased genetic risk, and that possibility has been raised for childhood onset schizophrenia [Rutter et al., 1999].

We have phenotyped 10 nuclear families with both the P50 paradigm and smooth pursuit eye movements. In contrast to families of adult onset probands, who almost always show unilineal presence of the P50 inhibitory deficit in the parents of the probands, i.e., only one parent affected [Waldo et al., 1994, Ross et al., 1998], the families of the childhood onset probands generally show bilineal presence of the P50 inhibitory deficit, i.e., both parents affected [$P < 0.006$; Ross et al., 1999]. A similar pattern is observed with one of the smooth pursuit eye movement deficits, an increase in the level of small leading saccades. In addition, the severity of the eye tracking deficit in the childhood onset-probands

was twice as great as that found in adult onset probands or in the children's parents. These findings thus support the hypothesis of possible co-dominant inheritance in childhood onset schizophrenia.

GENOCOPIES OF THE LINKAGE AT THE $\alpha 7$ NICOTINIC RECEPTOR LOCUS

Genocopies for schizophrenia are not common, but there have been reports of schizophrenic like illnesses near the chromosome 15q14 $\alpha 7$ nicotinic receptor locus, associated with a variety of genetic abnormalities. For example, a recent cytogenetic investigation implicates the chromosome 15q14 region in two cases of schizophrenia in an extended pedigree [Calzolari et al., 1996]. In this family, meioses involving a balanced translocation with a breakpoint at 15q13–14 produced a satellite chromosome in two cousins whose mothers both carried the translocation. Both individuals developed schizophrenia, that

These findings thus support the hypothesis of possible co-dominant inheritance in childhood onset schizophrenia.

was otherwise not present in the family's history. Three other genetic illnesses linked to this chromosomal region have been reported to co-express psychoses resembling schizophrenia in some individuals: Prader–Willi syndrome, a disease involving deletion or abnormal imprinting of genes at 15q11–12, Marfan syndrome that generally involves mutation of the fibrillin gene at 15q21, and Andersmann agenesis of the corpus callosum, that is caused by a deletion at chromosome 15q15 [Sirota et al., 1990; Clarke, 1993; Casaubon et al., 1996]. Either cosegregation of nearby genetic defects on

chromosome 15 or a common defect that affects expression of several genes in this region could be responsible for co-expression of schizophrenia with these illnesses.

SUMMARY

Initial neurobiological and genetic evidence of inheritance of a physiological abnormality related to schizophrenia has been obtained, that implicates the $\alpha 7$ nicotinic receptor gene. Bilineal inheritance may be associated with more severe, earlier onset illness. Replication of these findings, identification of the molecular abnormality in the gene, and determination of the full role of the abnormality in psychotic illnesses are necessary next steps to determine the role of this gene in the pathogenesis of schizophrenia.

REFERENCES

- Adams CE, Stevens KE. 1998. Inhibition of nitric oxide synthetase disrupts inhibitory gating of auditory responses in rat hippocampus. *J Pharmacol Exp Ther* 287:760–765.
- Adler LE, Pachtman E, Franks R, Pecevic M, Waldo MC, Freedman R. 1982. Neurophysiological evidence for a deficit in inhibitory mechanisms involved in sensory gating in schizophrenia. *Biol Psychiatry* 17: 639–654.
- Adler LE, Gerhardt GA, Franks R, Baker N, Nagamoto H, Drebing C, Freedman R. 1989. Sensory physiology and catecholamines in schizophrenia and mania. *Psychiatr Res* 31:297–309.
- Adler LE, Hoffer LJ, Griffith J, Waldo M, Freedman R. 1992. Normalization by nicotine of deficient auditory sensory gating in the relatives of schizophrenics. *Biol Psychiatry* 32: 607–616.
- Adler LE, Hoffer LD, Wiser A, Freedman R. 1993. Normalization of auditory physiology by cigarette smoking in schizophrenic patients. *Am J Psychiatr* 150:1856–1861.
- Andreasen NC, Paradiso S, O'Leary DS. 1998. "Cognitive dysmetria" as an integrative theory of schizophrenia: a dysfunction in cortical subcortical-cerebellar circuitry? *Schizophr Bull* 24:203–218.
- Arolt V, Lencer R, Nolte A, Muller-Myhsok B, Purnmann S, Schurmann M, Leutelt J, Pinnow M, Schwinger E. 1996. Eye tracking dysfunction is a putative phenotypic susceptibility marker of schizophrenia and maps to a locus on chromosome 6p in families with multiple occurrence of the disease. *Am J Med Genet* 67:564–579.
- Baker NJ, Staunton M, Adler LE, Gerhardt GA, Drebing C, Waldo M, Nagamoto H, Freedman R. 1990. Sensory gating deficits in psychiatric inpatients: Relation to catecholamine metabolites in different diagnostic groups. *Biol Psychiatry* 27:519–528.
- Berrettini WH, Ferraro TN, Goldin LR, Detera-Wadleigh SD, Choi H, Munie D, Guroff JJ, Kazuba DM, Nurnberger JI Jr, Hsieh WT, Hoehe MR, Gershon ES. 1997. A linkage study of bipolar illness. *Arch Gen Psychiatry* 54:27–35.
- Bickford PC, Luntz-Leybman V, Freedman R. 1993. Auditory sensory gating in the rat hippocampus: modulation by brainstem activity. *Brain Res* 607:33–38.
- Bickford-Wimer PC, Nagamoto H, Johnson R, Adler LE, Egan M, Rose GM, Freedman R. 1990. Auditory sensory gating in hippocampal neurons: a model system in the rat. *Biol Psychiatry* 27:183–192.
- Bickford PC, Wear KD. 1995. Restoration of sensory gating of auditory evoked response by nicotine in fimbria-fornix lesioned rats. *Brain Res* 705:235–240.
- Blouin JL, Dombroski BA, Nath SK, Lasseter VK, Wolyniec PS, Nestadt G, Thorndike M, Ullrich G, Mcgrath J, Kasch L, Lamacz M, Thomas MG, Gehrig C, Radhakrishna U, Snyder SE, Balk KG, Neufeld K, Swartz KL, Demarchi N, Papadimitriou GN, Dikeos DG, Stefanis CN, Chakravarti A, Childs B, Housman DE, Pulver AE. 1998. Schizophrenia susceptibility loci on chromosome 13q31 and 8p21. *Nature Genet* 20: 70–73.
- Borecki IB, Rao DC, Yaouanq J, Lalouel JM. 1990. Serum ferritin as a marker of affection for genetic hemochromatosis. *Human Hered* 40:159–166.
- Boutros NN, Zouridakis G, Overall J. 1991. Replication and extension of P50 findings in schizophrenia. *Clin Electroencephalogr* 22:40–45.
- Breese CR, Adams C, Logel J, Drebing C, Rollins Y, Barnhart M, Sullivan B, DeMasters BK, Freedman R, Leonard S. 1997. Comparison of the regional expression of nicotinic acetylcholine receptor $\alpha 7$ mRNA and [125 I]- α -bungarotoxin binding in human postmortem brain. *J Comp Neurol* 387: 385–398.
- Calzolari E, Aiello V, Pallazzi P, Sensi A, Calzolari S, Orrico D, Calliari L, Holler H, Marzi C, Belli S, Bernardi F, Patracchini P. 1996. Psychiatric disorder in a familial 15;18 translocation and sublocalization of myelin basic protein on 18q22.3. *Am J Med Genet* 67:154–161.
- Cannon TD, Zorilla LE, Shatslef LE. 1994. Neuropsychological functioning in siblings discordant for schizophrenia and healthy volunteers. *Arch Gen Psychiatry* 51:651–661.
- Cao Q, Martinez M, Zhang J, Sanders AR, Badner JA, Cravchik A, Markey C, Beshah E, Guroff JJ, Maxwell ME, Kazuba D, Goldin LR, Gershon EV, Gejman P. 1997. Suggestive evidence for a schizophrenia susceptibility locus on chromosome 6q and a confirmation in an independent series of pedigrees. *Genomics* 43:1–8.
- Casaubon LK, Melanson M, Lopes-Cendes I, Marineau C, Andersmann E, Andersmann F, Weissenbach J, Prevost C, Bouchard JP, Mathieu J, Rouleau GA. 1996. The gene responsible for a severe form of peripheral neuropathy and agenesis of the corpus callosum maps to chromosome 15q. *Am J Hum Genet* 58:28–34.

- Chini B, Raimond E, Elgoyhen AB, Moralli D, Balzaretto M, Heinemann S. 1994. Molecular cloning and chromosomal localization of human $\alpha 7$ -nicotinic receptor subunit gene (CHRNA7). *Genomics* 19:379–381.
- Clarke DJ. 1993. Prader-Willi syndrome and psychoses. *Br J Psychiatry* 163:680–684.
- Clementz BA, Geyer MA, Braff DL. 1997. P50 suppression among the relatives of schizophrenia patients. *Schizophr Res* 24:232.
- Clementz BA, Geyer MA, Braff DL. 1998. Multiple site evaluation of P50 suppression among schizophrenia and normal comparison subjects. *Schizophr Res* 30:71–80.
- Coon H, Plaetke R, Holik J, Hoff M, Myles-Worsley M, Freedman R, Byerley W. 1993. Use of a neurophysiological trait in linkage analysis of schizophrenia. *Biol Psychiatry* 34:277–289.
- Cornblatt BA, Keilp JG. 1994. Impaired attention, genetics, and the pathophysiology of schizophrenia. *Schizophr Bull* 20:31–46.
- Cullum CM, Harris JG, Waldo MC, Smernoff E, Madison A, Nagamoto HT, Griffith J, Adler LE, Freedman R. 1993. Neurophysiological and neuropsychological evidence for attentional dysfunction in schizophrenia. *Schizophr Res* 10:131–141.
- Curtis L, Blouin JL, Radhakrishna U, Gehring C, Lasseter VK, Wolyniec P, Dombroski B, Kazazian HH, Pulver AE, Housman D, Bertrand D, Antonarakis SE. 1999. No evidence of linkage between schizophrenia and markers at chromosome 15q13–14. *Am J Med Genet* 88:109–112.
- de Amicis LA, Wagstaff DA, Cromwell RL. 1986. Reaction time crossover as a marker of schizophrenia and of higher functioning. *J Nerv Ment Dis* 174:177–179.
- Edenberg HJ, Foroud T, Conneally PM, Sorbel JJ, Carr K, Crose C, Willig C, Zhao J, Miller M, Bowman E, Mayeda A, Rau NL, Smiley C, Rice JP, Goate A, Reich T, Stine OC, McMahon F, DePaulo JR, Meyers D, Detera-Wadleigh SD, Goldin LR, Gershon ES, Blehar MC, Nurnberger Jr JL. 1997. Initial genomic scan of the NIMH genetics initiative bipolar pedigrees: chromosomes 3, 5, 15, 16, 17, and 22. *Am J Med Genet* 74:238–246.
- Erwin RJ, Mawhinney-Hee M, Gur RC, Gur RE. 1991. Mid-latency auditory evoked responses in schizophrenia. *Biol Psychiatry* 30:430–442.
- Erwin RJ, Turetsky BJ, Moberg P, Gur RC, Gur RE. 1998. P50 abnormalities in schizophrenia: relationship to clinical and neurophysiological indices of attention. *Schizophr Res* 33:157–167.
- Frazier CJ, Rollins YD, Breese CR, Leonard S, Freedman R, Dunwiddie TV. 1997. Acetylcholine activates an α -bungarotoxin sensitive nicotinic current in rat hippocampal interneurons, but not pyramidal cells. *J Neurosci* 18:1187–1195.
- Freedman LR, Rock D, Roberts SA, Cornblatt BA, Erlenmeyer-Kimling L. 1998. The New York High-Risk Project: attention, anhedonia and social outcome. *Schizophr Res* 30:1–9.
- Freedman R, Adler LE, Bickford P, Byerley W, Coon H, Cullum MC, Griffith JM, Harris JG, Leonard S, Miller C, Myles-Worsley M, Nagamoto HT, Rose GM, Waldo M. 1994. Schizophrenia and nicotinic receptors. *Harvard Rev Psychiatry* 2:179–192.
- Freedman R, Coon H, Myles-Worsley M, Orr-Urtreger A, Olincy A, Davis A, Polymeroopoulos M, Holik J, Hopkins J, Hoff M, Rosenthal J, Waldo MC, Reimherr F, Wender P, Yaw J, Young DA, Breese CR, Adams C, Patterson D, Adler LE, Kruglyak L, Leonard S, Byerley W. 1997. Linkage of a neurophysiological deficit in schizophrenia to a chromosome 15 locus. *Proc Natl Acad Sci* 94:587–592.
- Freedman R, Wetmore C, Stromberg I, Leonard S, Olson L. 1993. α -Bungarotoxin binding to hippocampal interneurons: immunocytochemical characterization and effects on growth factor expression. *J Neurosci* 13:1965–1975.
- Gault J, Robinson M, Berger R, Drebing C, Logel J, Hopkins J, Moore T, Jacobs S, Merriwether J, Choi MJ, Kim EJ, Walton K, Buiting K, Davis A, Breese C, Freedman R, Leonard S. 1998. Genomic organization and partial duplication of the human $\alpha 7$ neuronal nicotinic acetylcholine receptor gene. *Genomics* 52:173–185.
- Goldberg TE, Rigland JD, Torrey EF, Gold JM, Bigelow LB, Weinberger DR. 1990. Neuropsychological assessment of monozygotic twins discordant for schizophrenia. *Arch Gen Psychiatry* 47:1066–1072.
- Griffith J, Hoffer LD, Adler LE, Zerbe GO, Freedman R. 1995. Effects of sound intensity on a mid-latency evoked response to repeated auditory stimuli in schizophrenic and normal subjects. *Psychophysiology* 32:460–466.
- Harris JG, Adler LE, Young DA, Cullum CM, Rilling LM, Cicerello A, Intemann PM, Freedman R. 1996. Neuropsychological dysfunction in parents of schizophrenics. *Schizophr Res* 20:253–260.
- Hershman KM, Freedman R, Bickford PC. 1995. GABA-B antagonists diminish the inhibitory gating of auditory response in the rat hippocampus. *Neurosci Lett* 190:133–136.
- Holzman PS, Kringlen E, Matthyse S, Flanagan SD, Lipton RB, Cramer G, Levin S, Lange K, Levy DL. 1988. A single dominant gene can account for eye tracking dysfunction and schizophrenia in offspring of discordant twins. *Arch Psychiatry* 45:641–647.
- Jin Y, Potkin SG, Patterson JV, Sandman CA, Hetrick WP, Bunney Jr WE. 1997. Effects of P50 temporal variability on sensory gating in schizophrenia. *Psychiatry Res* 70:71–81.
- Judd LL, McAdams L, Budnick B, Braff DL. 1992. Sensory gating deficit in schizophrenia: new results. *Am J Psychiatry* 149:488–493.
- Kaufmann CA, Suarez B, Malaspina D, Pepple J, Svrakic D, Markel PD, Meyer J, Zambuto CT, Schmitt K, Matisse TC, Friedman JMH, Hampe C, Lee H, Shore D, Wynne D, Faraone SV, Tsuang MT, Cloninger CR. 1998. NIMH Genetics Initiative Millennium Consortium—linkage analysis of African-American pedigrees. *Am J Med Genet* 81:282–289.
- Leonard S, Gault J, Moore T, Hopkins J, Robinson M, Olincy A, Adler LE, Cloninger R, Kaufmann CA, Tsuang MT, Faraone SV, Malaspina D, Svrakic DM, Freedman R. 1998. Further investigation of a chromosome 15 locus in schizophrenia: analysis of affected sib pairs from the NIMH Genetics Initiative. *Neuropsychiatr Genet* 81:308–312.
- Leppert M, Burt R, Hughes JP, Samowitz W, Nakamura Y, Woodward S, Gardner E, Lalouel JM, White R. 1990. Genetic analysis of an inherited predisposition to colon cancer in a family with a variable number of adenomatous polyps. *N Engl J Med* 322:904–908.
- Luntz-Leyman V, Bickford PC, Freedman R. 1992. Cholinergic gating of response to auditory stimuli in rat hippocampus. *Brain Res* 587:130–136.
- McKenna K, Gordon CT, Rapoport JL. 1994. Childhood onset schizophrenia: timely neurobiological research. *J Am Acad Child Adol Psychiatry* 33:771–781.
- Miller CL, Freedman R. 1993. Medial septal neuron activity in relation to an auditory sensory gating paradigm. *Neuroscience* 55:373–380.
- Miller CC, Freedman R. 1995. The activity of hippocampal interneurons and pyramidal cells during the response of the hippocampus to repeated auditory stimuli. *Neuroscience* 69:371–381.
- Nagamoto HT, Adler LE, Waldo MC, Freedman R. 1989. Sensory gating in schizophrenics and normal controls: effects of changing stimulation interval. *Biol Psychiatry* 25:549–561.
- Nelson MD, Saykin AJ, Flashman LA, Riordan HJ. 1998. Hippocampal volume reduction in schizophrenia as assessed by magnetic resonance imaging: a meta-analytic study. *Arch Gen Psychiatry* 55:433–440.
- Neves-Pereira M, Braganza-King N, Walker ML, Kennedy JL. 1999. TDT studies of CHRNA7 in schizophrenia and bipolar disorder. *Schizophr Res* 36:93.
- Neves-Pereira M, Bassett AS, Honer WG, Lang D, King NA, Kennedy JL. 1998. No evidence for linkage of the CHRNA7 gene region in Canadian schizophrenia families. *Am J Med Genet* 81:361–363.
- Olincy A, Ross RG, Young DA, Roath M, Freedman R. 1998. Improvement in smooth pursuit eye movements after cigarette smoking in schizophrenic patients. *Neuropsychopharmacology* 18:175–185.
- Olincy A, Young DA, Freedman R. 1997. Increased levels of nicotine metabolite cotinine in schizophrenic smokers compared to other smokers. *Biol Psychiatry* 42:1–5.
- Ross RG, Harris JG, Olincy A, Radant A, Adler LE, Freedman R. 1998. Anticipatory saccades during smooth pursuit eye movements and familial transmission of schizophrenia. *Biol Psychiatry* 8:690–697.
- Ross RG, Olincy A, Harris JG, Radant A, Hawkins M, Adler LE, Freedman R. 1999. Evidence for bilineal inheritance of physiological indicators of risk in childhood-onset schizophrenia. *Am J Med Genet Neuropsychiatr Genet* 88:188–199.
- Rutter M, Silberg J, O'Connor T, Simonoff E. 1999. Genetics and child psychiatry: II Empirical research findings. *J Child Psychol Psychiatry* 40:19–55.
- Sham PC, Morton NE, Muir WJ, Walker M, Collins A, Shields DC, St Clair DM, Blackwood DH. 1994. Segregation analysis of

- complex phenotypes: an application to schizophrenia and auditory P300 latency. *Psychiatr Genet* 4:29–38.
- Shihabuddin L, Silverman JM, Buchsbaum MS, Seiver LJ, Luu C, Germans MK, Metzger M, Mohs RC, Smith CJ, Spiegel-Cohen J, Davis KL. 1996. Ventricular enlargement associated with linkage marker for schizophrenia-related disorders in one pedigree. *Mol Psychiatry* 1:215–222.
- Sirota P, Frydman M, Sirota L. 1990. Schizophrenia and Marfan syndrome. *Br J Psychiatry* 157:433–436.
- Stevens KE, Freedman R, Collins AC, Hall M, Leonard S, Marks MJ, Rose GM. 1996. Genetic correlation of inhibitory gating of hippocampal auditory evoked response and alpha-bungarotoxin-binding nicotinic cholinergic receptors in inbred mouse strains. *Neuropsychopharmacology* 15:152–162.
- Tamminga CA, Thaker GK, Buchanan R, Kirkpatrick B, Alphas LD, Chase TN, Carpenter WT. 1992. Limbic system abnormalities identified in schizophrenia using positron emission tomography with fluorodeoxyglucose and neocortical alterations with deficit syndrome. *Arch Gen Psychiatry* 49:522–530.
- Tsuang MT, Lyons MJ, Faraone SV. 1990. Heterogeneity of schizophrenia: conceptual models and analytic strategies. *Br J Psychiatry* 156:17–26.
- Waldo MC, Cawthra E, Adler LE, Dubester S, Staunton M, Nagamoto H, Baker N, Madison A, Simon J, Scherzinger A, Drebing C, Gerhardt G, Freedman R. 1994. Auditory sensory gating, hippocampal volume, and catecholamine metabolism in schizophrenics and their siblings. *Schizophr Res* 12:93–106.
- Waldo M, Myles-Worsley M, Madison A, Byerley W, Freedman R. 1995. Sensory gating deficits in parents of schizophrenics. *Am J Med Genet* 60:506–511.
- Ward PB, Hoffer LD, Liebert BJ, Catts SV, O'Donnell M, Adler LE. 1996. Replication of a P50 auditory gating deficit in Australian patients with schizophrenia. *Psychiatry Res* 64:121–135.
- Yee CM, Neuchterlain KH, Morris SE, White PM. 1998. P50 suppression in recent-onset schizophrenics: clinical correlates and risperidone effect. *J Abnorm Psych* 107:691–698.